

John Peters, PhD

Assoc. Director, Outcomes & Technology Regional Mental Health & Chemical Dependency Kaiser Permenente, Northern California



From Depression Monitoring to a Transdiagnostic Approach

- Priorities: Clinical outcomes, Quality improvement, Research
- Aligning behavioral health care across primary and specialty care
- Shifting focus to implementation: Transdiagnostic tool found to be most feasible in Mental Health
- Developed AOQ with PHQ9, GAD2, 2 functioning items
 - + alliance items in MH settings




2.3 years of implementation of the AOQ

- AOQ global distress scores in outpatient Mental Health:
 - In 2013, through Q3: 270,000 global distress scores
 - 70% of all adult MD/NP visits
 - 60% of all adult individual psychotherapy visits
 - 25% of all adult group psychotherapy visits
 - 54% combined




Progress Data Drives MH Implementation

Measurement & Monitoring



Progress Data Drives MH Implementation

Measurement & Monitoring



Overall improvement of outcomes

Understand outcomes of specific programs


Identify bright spots in the program

New knowledge about care modalities and pathways





Overall improvement of outcomes: Feedback improves outcomes

KP NCAL pilot (3 sites)



A non-KP PPO network





3. Identify bright spots in the program

Medical Sites	N	Severity- Adjusted Effect Size
1	542	0.77
2	385	0.89
3	858	0.91
4	944	0.91
5	1,707	0.95
6	1,462	1.00
7	1,570	1.01
8	625	1.04
9	423	1.08
10	376	1.35






4. New knowledge about internal care modalities and pathways, ex. 1

Effect size of predominant treatment modalities

2800 cases with 3 or more assessments at 3 clinics





4. New knowledge about internal care modalities and pathways, ex. 2

Impact of assessing "Goodness of Fit"

2400 cases with no GoF scores, vs. repeated Gof scores

