
GGC4H: Testing Feasibility and Effectiveness of Universal Parent-Focused Prevention in Three Healthcare Systems


Arne L. Beck, PhD Senior Investigator Institute for Health Research Kaiser Permanente






1








Study design Randomly assigned 75 pediatricians within 3 healthcare systems and 10 clinics Recruited ~1975 adolescents to the study – 2 cohorts Offered GGC to 512 enrolled parents in intervention arm RE-AIM* measurement framework - Implementation: Reach, adoption, implementation fidelity, participant engagement and skills - Effectiveness: Evaluate GGC's impact on adolescent health NIH PRAGMATIC TRIALS COLLABORATORY Rethinking Clinical Trials







7

EHR did not have the outcomes data GGC4H needed. *We developed a* Youth Behavioral Health Survey instead:

Primary	Secondary	Exploratory	Mechanisms to
Outcomes	Outcomes	Outcomes	Impact
Substance Use Age of Initiation Substances Examined Alcohol, Marijuana, Cigarettes, E-Cigarettes, Inhalants, Opioids, Other Drugs	Mental Health Depression (PHQ-9) Antisocial Behavior Ever Past-Year Substance Use Lifetime Frequency Past-Year, Past 30-day Use Past 30-day Use Amount	Anxiety (GAD-7) Screen & Social Media Time Sexting	Parent and Family Risk & Protective Factors (RPFs) Individual RPFs Peer RPFs School RPFs

- Developed Adolescent Behavioral Health Survey to collect data on behavioral health outcomes; widely used, validated measures
- Administered online or by telephone with trained interviewers

Rethinking Clinical Trials®

